

Catabolism of proteins, fats, carbohydrates in 3 stages of cellular respiration

Stage 1: oxidation of fatty acids, glucose, and some amino acids yields acetyl-CoA.

Stage 2: oxidation of acetyl groups in the citric acid cycle includes four steps in which electrons are abstracted.

Stage 3: electrons carried by NADH and FADH₂ are funneled into a chain of mitochondrial (or, in bacteria, plasma membrane-bound) electron carriers—the respiratory chain—ultimately reducing O_2 to H₂O. This electron flow drives the production of ATP.

Carbohydrates

- Polysaccharides most abundant CARBS in diet
- Starch & cellulose By plant food
- Glycogen by food of animal origin
- Starch & glycogen digestion starts in mouth
- Salivary amylase hydrolyze alpha 1-4 glycosidic linkages
- Yield a mixture of maltose, glucose & oligosaccharides
- Digestion stops in stomach
- Continue in small intestine By pancreatic amylase
- Cellulose cannot be digested
- Lack of enzyme to hydrolyze beta 1-4 linkage
- Undigested cellulose give bulk of fiber (roughage)
- This is desirable for proper motility of intestine
- Disaccharides digestion in small intestine
- Sucrose by sucrase (invertase), Lactose by lactase (β-galactosidase), Maltose by maltase

CARBOHYDRATE METABOLISM Glycolysis

- D-glucose a major fuel, occupies a central position in metabolism, relatively rich in potential energy
 (complete oxidation to carbon dioxide and water proceeds with a standard free energy of -2,840 kJ/mol)
- Glucose a precursor, provides metabolic intermediates, the necessary starting material for biosynthetic reactions

In higher plants and animals, glucose has four major fates



CARBOHYDRATE METABOLISM Glycolysis

- Glycolysis (Greek glykys, meaning "sweet" and lysis, meaning "splitting")
- Described by Gustav Embden, Otto Meyerhof and Jacob Parnas
- Known as Embden-Meyerhof-Parnas pathway (EMP)
- Reactions of Glycolysis take place in the cytosol of cells
- Unique; it can utilize O_2 if available and can also work in the absence of O_2
- Glycolytic sequence of reactions differ from one specie to another only in:

1. how the rate is regulated 2. metabolic fate of pyruvate

- Breakdown of the (6C) 1 glucose into 2 molecules of the (3C) pyruvate occurs in 10 steps (reactions)
- First 5 reactions Phase 1 (Preparatory phase/energy investment phase)
- Last 5 reactions Phase 2 (Payoff Phase/energy production phase/oxidative phase)



Figure 14-2a *Lehninger Principles of Biochemistry, Fifth Edition* © 2008 W.H. Freeman and Company



For each molecule of glucose that passes through preparatory phase, 2 molecules of glyceraldehyde-3phosphate are formed; both pass through the payoff phase.

Pyruvate is the end product of second phase of glycolysis.

For each alucose molecule. are consumed the in bhase preparatory are broduced in the bavoti bhase. a net aivina ber molecule alucose converted to pyruvate.

3 Important Chemical Transformations

- 1. Degradation of C skeleton of glucose to yield pyruvate
- 2. Phosphorylation of ADP to ATP
- 3. Transfer of H atoms or e to NAD+, forming NADH

Overall Equation for Glycolysis

 $\begin{array}{l} Glucose + 2NAD^{+} + 2ADP + 2P_{i} \longrightarrow \\ 2 \ pyruvate + 2NADH + 2H^{+} + 2ATP + 2H_{2}O \end{array}$



this is calculated as 2.5ATP per NADH



Three possible catabolic fates of pyruvate formed in glycolysis

Lactic acid fermentation

- Vigorously contracting skeletal muscle function anaerobically, pyruvate cannot be oxidized further due to lack of oxygen. So, pyruvate is reduced to lactate
- Certain tissues & cells (retina, brain, RBCs) convert glucose to lactate even under aerobic conditions (as these don't have mitochondria)
- Lactate (the dissociated form of lactic acid) is also the product of glycolysis under anaerobic conditions in microorganisms that carry out the lactic acid fermentation



Ethanol or Alcohol fermentation

In some plant tissues & in certain invertebrates, protests & microorganisms such as brewer's yeast, pyruvate is converted anaerobically into ethanol & CO_2





David L. Nelson and Michael M. Cox

LEHNINGER PRINCIPLES OF BIOCHEMISTRY Fifth Edition



CHAPTER 16 The Citric Acid Cycle

Hans Krebs, 1900-1981

Unnumbered 16 p615 Lehninger Principles of Biochemistry, Fifth Edition < 2008 W.H.Freeman and Company

© 2008 W. H. Freeman and Company





 2 C atoms enter the cycle as acetyl CoA & leave as CO₂

Energy produced 3 NAD+ reduced to 3NADH=+7.5 * ATPs 1 FAD reduced to 1FADH2 =+1.5 * ATPs <u>1 GDP converted to GTP =+1 ATP</u> Net = 10 ATPs for 1 acetyl-CoA

FOR 2 acetyl-CoA=20 ATPs

*this is calculated as 2.5ATP per NADH and 1.5ATP per FADH₂



Products of one turn of the citric acid cycle

At each turn of the cycle, 3NADH, 1 FADH₂, 1 GTP (or ATP), and 2 CO_2 are released in oxidative decarboxylation reactions.

Here all cycle reactions are shown as proceeding in one direction only, but keep in mind that most of the reactions are reversible