

Biochem 717

Gene Expression

Prof Amer Jamil

Dept. of Biochemistry

University of Agriculture

Faisalabad

4.3 Methods of Expressing Cloned Genes

Cloning a gene permits

- Production of large quantities of a particular DNA sequence for detailed study
- Large quantities of the gene's product can also be obtained for further use
 - Study
 - Commerce

Expression Vectors

- Vectors discussed so far are used to first put a foreign DNA into a bacterium to replicate and screen
- **Expression vectors** are those that can yield protein products of the cloned genes
 - Bacterial expression vectors typically have two elements required for active gene expression; a strong promoter and a ribosome binding site near an initiating codon






Figure 9-10
Lehninger Principles of Biochemistry, Fifth Edition
 © 2008 W.H. Freeman and Company

Fusion Proteins

- Some cloning vectors, pUC and pBS, can work as expression vectors using *lac* promoter
- If inserted DNA is in the same reading frame as interrupted gene, a fusion protein results
 - These have a partial β -galactosidase sequence at amino end
 - Inserted cDNA protein sequence at carboxyl end





Glutathione-S-transferase (GST)

Figure 9-12a

Lehninger Principles of Biochemistry, Fifth Edition

© 2008 W. H. Freeman and Company




Figure 9-12b

Lehninger Principles of Biochemistry, Fifth Edition

© 2008 W. H. Freeman and Company




Figure 9-12b part 1

Lehninger Principles of Biochemistry, Fifth Edition

© 2008 W. H. Freeman and Company




Figure 9-12b part 2
Lehninger Principles of Biochemistry, Fifth Edition
© 2008 W. H. Freeman and Company

Inducible Expression Vectors

- Main function of expression vector is to yield the product of a gene – usually more is better
- For this reason, expression vectors have very strong promoters
- It is usually advantageous to keep a cloned gene repressed until time to express
 - Large quantities of eukaryotic protein in bacteria are usually toxic
 - Can accumulate to levels that interfere with bacterial growth
 - Expressed protein may form insoluble aggregates, called inclusion bodies

Controlling the *lac* Promoter

- *lac* promoter is somewhat inducible
 - Stays off until stimulated by inducer IPTG
 - However, repression is typically incomplete or leaky and some expression will still occur
- To avoid this problem, use a plasmid or phagemid carrying its own *lacI* repressor gene to keep the cloned gene off until it is induced by IPTG

Alternatives to the *lac* Promoter

- The hybrid *trc* promoter combines the strength of the *trp* (tryptophan operon) promoter with the inducibility of the *lac* promoter
- Promoter from *ara* operon, P_{BAD} , allow fine control of transcription
 - Inducible by arabinose, a sugar
 - Transcription rate varies with arabinose concentration

Alternatives to the *lac* Promoter


- The lambda (λ) phage promoter, P_L , is tightly controlled
- Expression vectors with this promoter-operator system are used in host cells with temperature-sensitive λ repressor gene
 - Repressor functions at low temperatures
 - Raise temperature above the nonpermissive level (42°C) and the repressor doesn't function and the cloned gene is expressed

Expression Vectors That Produce Fusion Proteins

- Most vectors express fusion proteins
 - The actual natural product of the gene isn't made
 - Extra amino acids help in purifying the protein product
- Oligohistidine expression vector has a short sequence just upstream of MCS encoding 6 His
 - Oligohistidine has a high affinity for divalent metal ions like nickel (Ni^{2+})
 - Permits purification by nickel affinity chromatography
 - The his tag can be removed using enzyme enterokinase without damage to the protein product

Using an Oligohistidine Expression Vector


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Expression vector λ gt11


- This phage contains the *lac* control region followed by the *lacZ* gene
- The cloning sites are located within the *lacZ* gene
- Products of gene correctly inserted will be fusion proteins with a β -galactosidase leader

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Detecting positive λ gt11 clones via antibody screening

- Lambda phages with cDNA inserts are plated
- Protein released are blotted onto a support
- Probe with antibody specific to protein
- Antibody bound to protein from plaque is detected with labeled protein A



Bacterial Expression System Shortcomings

- There are problems with expression of eukaryotic proteins in a bacterial system
 - Bacteria may recognize the proteins as foreign and destroy them
 - Post-translational modifications are different in bacteria
 - Bacterial environment may not permit correct protein folding
- Very high levels of cloned eukaryotic proteins can be expressed in useless, insoluble form

Eukaryotic Expression Systems

- Avoid bacterial expression problems by expressing the protein in a eukaryotic cell
- Initial cloning done in *E. coli* using a **shuttle vector**, able to replicate in both bacterial and eukaryotic cells
- Yeast is suited for this purpose
 - Rapid growth and ease of culture
 - A eukaryote with more appropriate post-translational modification
 - Use of the yeast export signal peptide secretes protein into growth medium for easy purification

Use of Baculovirus As Expression Vector

- Viruses in this class have a large circular DNA genome, 130 kb
- Major viral structural protein is made in huge amounts in infected cells
 - The promoter for this protein, polyhedrin, is very active
 - These vectors can produce up to 0.5 g of protein per liter of medium
 - Nonrecombinant viral DNA entering cells does not result in infectious virus as it lacks an essential gene supplied by the vector

Expressing a Gene in a Baculovirus

Fig. 4.20a




Fig. 4.20b

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



© Dr. Robert Turgeon and Dr. B. Gillian Turgeon, Cornell University.

Animal Cell Transfection

- Carried out in two ways:
- Calcium phosphate
 - Mix cells with DNA in a phosphate buffer and add a solution of calcium salt to form a precipitate
 - The cells take up the calcium phosphate crystals, which include some DNA
- Liposomes
 - The DNA is mixed with lipid to form liposomes, small vesicles with some of the DNA inside
 - DNA-bearing liposomes fuse with the cell membrane to deliver DNA inside the cell

Using the Ti Plasmid to Transfer Genes to Plants

- Genes can be introduced into plants with vectors that can replicate in plant cells
- Common bacterial vector promoters and replication origins are not recognized by plant cells
- Plasmids are used containing T-DNA
 - T-DNA is derived from a plasmid known as tumor-inducing (Ti)
 - Ti plasmid comes from bacteria that cause plant tumors called crown galls


Ti Plasmid Infection

- Bacterium infects plant, transfers Ti plasmid to host cells
- T-DNA integrates into the plant DNA causing abnormal proliferation of plant cells
- T-DNA genes direct the synthesis of unusual organic acids, opines which can serve as an energy source to the infecting bacteria but are useless to the plant

The Ti Plasmid Transfers Crown Gall


Fig. 4.21

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Use of the T-DNA Plasmid

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Reporter Gene:

A gene with a readily measurable phenotype that can be easily distinguished over a background of endogenous proteins

OR

A gene that researchers attach to a regulatory sequence of another gene of interest in cell culture, animals or plants.

Reporter Assay

1. Measures gene expression or transcriptional activity
2. Assay of transcription factors.
3. DNA promoter assay
4. Confirmation of transgenesis


Reporter Assays

1. CAT: chloramphenicol acetyltransferase
2. β -gal (β -galactosidase):
3. **GUS Reporter Gene System**
4. **SEAP (secreted alkaline phosphatase):**
5. Luciferase
6. GFP: Green Fluorescent Protein
7. **Bioluminescence Resonance Energy Transfer (BRET)**
8. **Human Growth Hormone (hGH) Reporter Gene System**


CAT: chloramphenicol acetyltransferase

1. 1st reporter gene used to monitor transcriptional activity in cells
2. Bacterial enzyme that transfers acetyl groups from acetyl-CoA to chloramphenicol, detoxifying it
3. Reaction quantified using radiolabeled substrates (^{14}C -chloramphenicol) or by ELISA (non-radioactive)

CAT assay: acetylated & non-acetylated chloramphenicol are checked by TLC





CAT assay: ELISA



β -gal (β -galactosidase):



- *E. coli* enzyme (encoded by *lacZ*) that hydrolyzes galactosidase sugars such as lactose
- Many assay formats: colorimetric, fluorescent, chemiluminescent



GUS Reporter Gene System

GUS encodes the beta-glucuronidase enzyme from *E. coli*.

An active enzyme may be detected using X-gal, which forms an intense blue product after cleavage by β -galactosidase



Luciferase:





Renilla reniformis



Photinus pyralis

- . Firefly (*Photinus pyralis*) luciferase
- . Sea pansy (*Renilla reniformis*) luciferase
- . Firefly luciferase produces light by ATP-dependent oxidation



- . Bioluminescence or light emission is determined by a luminometer

Dual-Luciferase[®] and Dual-Luciferase[®] 1000 Assay Protocols

Assay with 96-Well Plate

Before you begin:


Set injectors 1 and 2 to dispense 100 μ l of **LAR II** and **Stop & Glo[®] Reagent**, respectively.

For measurements, use a 1- to 2-second delay and a 5- to 10-second read time.

(inside luminometer)




Dispense 100 μ l of **LAR II**.



Measure firefly luciferase activity.

Dispense 100 μ l of **Stop & Glo[®] Reagent**.



Measure *Renilla* luciferase activity.

Repeat cycle for remaining wells in plate.

2864MA02_0A

Assay with Manual or Single-Injector Luminometer



Predispense 100 μ l of **LAR II** into luminometer tube.



Program luminometer.



Transfer 20 μ l of PLB Lysate. Mix.



Measure firefly luciferase activity.



Dispense 100 μ l of **Stop & Glo[®] Reagent**.





Measure *Renilla* luciferase activity.

2865MA02_0A

GFP: Green Fluorescent Protein

- Derived from jellyfish *Aequorea victoria*
- Autofluorescent upon UV irradiation (doesn't require cofactors or substrates)
- Retains activity in presence of heat, denaturants, detergents, most proteases
- Allows for non-invasive monitoring of gene expression in living tissues



SEAP (secreted alkaline phosphatase):

- Secreted outside the cell (can assay sample repeatedly and non-destructively by sampling culture medium)
- This protein is quantified directly by measuring the enzyme activity in the supernatant of the culture medium.
- Fluorescence and chemiluminescence assays are available for detection.




Bioluminescence Resonance Energy Transfer (BRET)

These systems are based on excitation energy transfer from a bioluminescent donor molecule, e.g. Renilla luciferase to a fluorescent acceptor molecule, like green fluorescent protein (GFP).

Renilla luciferase as well as GFP are both attached to the proteins to be examined.

In the presence of coelenterazine, a BRET signal is generated when the fusion proteins are associated.

Excellent discrimination of the *luc* and GFP signals results when coelenterazine 400a is used as the *luc* substrate.




Human Growth Hormone (hGH) Reporter Gene System

The human growth hormone (hGH) encoded reporter protein is secreted into the culture medium by transfected cells.

The hGH from the supernatant of the culture medium binds to the antibody on the plate.



Subsequently, the bound hGH is detected in two steps via a digoxigenin-coupled anti-hGH antibody and a peroxidase-coupled anti-digoxigenin antibody.

Bound peroxidase is quantified by incubation with a peroxidase substrate such as TMB (3,3',5,5'-tetramethylbenzidine)




Highly parallel DNA detection I

A) Spotted Arrays





Rockefeller



Highly parallel DNA detection II

B) On-Chip Synthesis




Affymetrix





Highly parallel DNA detection III

D) Quantitative PCR (qPCR)




Quantitative polymerase chain reaction (Q-PCR) is a modification of polymerase chain reaction used to rapidly measure the quantity of a product of polymerase chain reaction. It is preferably done in real-time, thus is an indirect method for quantitatively measuring starting amounts of DNA, complementary DNA or ribonucleic acid (RNA). This is commonly used for the purpose of determining whether a genetic sequence is present or not, and if it is present the number of copies in the sample. (en.wikipedia.org)

Online Detection of Product



(c) Roche

TaqMan Probes in Action!




S1 Mapping

Use S1 mapping to locate the ends of RNAs and to determine the amount of a given RNA in cells at a given time

- Label a ssDNA probe that can only hybridize to transcript of interest
- Probe must span the sequence start to finish
- After hybridization, treat with S1 nuclease which degrades ssDNA and RNA
- Transcript protects part of the probe from degradation
- Size of protected area can be measured by gel electrophoresis


S1 Mapping the 5' End

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



S1 Mapping the 3' End

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Summary

- In S1 mapping, a labeled DNA probe is used to detect 5'- or 3'-end of a transcript
- Hybridization of the probe to the transcript protects a portion of the probe from digestion by S1 nuclease, specific for single-stranded polynucleotides
- Length of the section of probe protected by the transcript locates the end of the transcript relative to the known location of an end of the probe
- Amount of probe protected is proportional to concentration of transcript, so S1 mapping can be quantitative
- RNase mapping uses an RNA probe and RNase

Transgenic Plants



**Round Up Ready
Soybeans are resistant to
herbicide**

Herbicide Tolerance, Insect
Resistance, quality traits

Soybean, Corn, Cotton, Canola
Tomato






**“Golden” rice with
beta-carotene and
extra iron**



**Bt Corn produces
its own pesticide**

Ti plasmid and YAC






The Gene Gun




PDS1000 Microparticle Delivery System



Genetically Modified Organisms (Transgenic)



- **Introducing or modifying specific genes to alter the phenotype of an organism**
- **Transgenic: organism that contains a gene from another species in all of its cells**
- **Transgenic Animals**
 - **Models of Human Disease**
 - **Produce Pharmaceuticals**



Dolly - 1996



Fertilization vs. Cloning (somatic cell nuclear transfer)



Cloning since Dolly

Cloning of this sort has now been done on cattle, pigs and mice also.

The success rate has improved considerably.



Guar: First cloned endangered species dies 2 days after birth

